



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Adams County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		85 %	(83% - 87%)	1,027
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: gray;">■</span> AVERAGE		92 %	(89% - 93%)	721
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: red;">▼</span> LOWER than average		66 %	(62% - 70%)	499
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: gray;">■</span> AVERAGE		76 %	(72% - 80%)	488
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: gray;">■</span> AVERAGE		77 %	(74% - 80%)	755
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> AVERAGE		88 %	(84% - 92%)	247
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		86 %	(81% - 90%)	218
Eye Exam	<span style="color: gray;">■</span> AVERAGE		53 %	(46% - 59%)	218
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> AVERAGE		76 %	(70% - 81%)	218
Kidney Disease Screening	<span style="color: gray;">■</span> AVERAGE		81 %	(75% - 85%)	218
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: red;">▼</span> LOWER than average		69 %	(67% - 71%)	2,178
Antidepressants	<span style="color: red;">▼</span> LOWER than average		81 %	(79% - 82%)	2,641
Antacid Medication (Proton Pump Inhibitors)	<span style="color: red;">▼</span> LOWER than average		81 %	(79% - 83%)	1,324

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

## LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average


Must have 160+ patients per measure to be publicly reported.

## Generic Prescription Drugs

Medication for ACE inhibitors or  
angiotensin II receptor blockers  
(ARBs) used to treat High Blood  
Pressure



**LOWER**  
than average




84 % (83% - 85%)

3,075

Medication for attention deficit  
hyperactivity disorder



**AVERAGE**



64 % (61% - 67%)


893

## Health Screenings

Screening for Breast Cancer - Ages  
52-69



**AVERAGE**




62 % (56% - 67%)

344

Screening for Colon Cancer



**LOWER**  
than average




40 % (35% - 46%)

315

Screening for Cervical Cancer



**LOWER**  
than average




63 % (60% - 66%)

977

Screening for Chlamydia



**AVERAGE**




52 % (45% - 58%)

213

Adolescent Well-Care Visits



**AVERAGE**



31 % (28% - 34%)

1,014

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Asotin County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		88 %	(85% - 91%)	560
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: gray;">■</span> AVERAGE		93 %	(91% - 95%)	486
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: red;">▼</span> LOWER than average		58 %	(53% - 63%)	338
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: red;">▼</span> LOWER than average		65 %	(60% - 70%)	303
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: red;">▼</span> LOWER than average		66 %	(61% - 70%)	422
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: red;">▼</span> LOWER than average		78 %	(71% - 83%)	174
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: gray;">■</span> AVERAGE		76 %	(75% - 78%)	2,016
Antidepressants	<span style="color: gray;">■</span> AVERAGE		90 %	(89% - 90%)	4,779
Antacid Medication (Proton Pump Inhibitors)	<span style="color: gray;">■</span> AVERAGE		89 %	(88% - 90%)	1,800
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	<span style="color: gray;">■</span> AVERAGE		89 %	(88% - 90%)	2,613
Medication for attention deficit hyperactivity disorder	<span style="color: gray;">■</span> AVERAGE		64 %	(62% - 66%)	2,038
<b>Health Screenings</b>					
Screening for Breast Cancer - Ages 52–69	<span style="color: gray;">■</span> AVERAGE		63 %	(56% - 69%)	203
Screening for Colon Cancer	<span style="color: gray;">■</span> AVERAGE		52 %	(46% - 59%)	218

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

## LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average


Must have 160+ patients per measure to be publicly reported.

## Health Screenings

Screening for Cervical Cancer



**AVERAGE**




**64 % (60% - 68%)**

**578**

Adolescent Well-Care Visits



**LOWER**  
than average



**20 % (17% - 23%)**

**597**

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP














**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Benton County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> <b>BETTER than average</b>		88 %	(88% - 89%)	10,385
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: green;">●</span> <b>BETTER than average</b>		95 %	(95% - 95%)	10,685
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(85% - 90%)	742
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: gray;">■</span> <b>AVERAGE</b>		90 %	(88% - 91%)	1,217
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		76 %	(75% - 77%)	5,960
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		77 %	(76% - 78%)	5,063
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		77 %	(76% - 78%)	7,590
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: red;">▼</span> <b>LOWER than average</b>		79 %	(77% - 81%)	1,717
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(86% - 90%)	1,079
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: red;">▼</span> <b>LOWER than average</b>		16 %	(13% - 20%)	532
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: green;">●</span> <b>BETTER than average</b>		90 %	(88% - 92%)	634
<b>Depression</b>					
Antidepressant Medication (12 Weeks)	<span style="color: gray;">■</span> <b>AVERAGE</b>		66 %	(61% - 71%)	363
Antidepressant Medication (6 Months)	<span style="color: gray;">■</span> <b>AVERAGE</b>		46 %	(41% - 51%)	363

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Chelan County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		84 %	(82% - 85%)	3,235
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: red;">▼</span> LOWER than average		90 %	(89% - 91%)	3,136
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: red;">▼</span> LOWER than average		84 %	(79% - 87%)	304
Child and Adolescent Access to Primary Care - Ages 12–24 Months	<span style="color: green;">●</span> BETTER than average		95 %	(93% - 97%)	444
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: green;">●</span> BETTER than average		80 %	(79% - 82%)	2,444
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: green;">●</span> BETTER than average		81 %	(79% - 82%)	2,154
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: green;">●</span> BETTER than average		79 %	(77% - 80%)	2,971
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> AVERAGE		92 %	(90% - 94%)	789
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> AVERAGE		90 %	(86% - 93%)	317
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> AVERAGE		83 %	(77% - 88%)	184
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		86 %	(83% - 89%)	496
Eye Exam	<span style="color: gray;">■</span> AVERAGE		54 %	(49% - 58%)	496
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> AVERAGE		73 %	(69% - 77%)	496
Kidney Disease Screening	<span style="color: red;">▼</span> LOWER than average		73 %	(69% - 77%)	496

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Clallam County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	▼ LOWER than average		77 %	(76% - 78%)	3,822
Adult Access to Preventive/Ambulatory Care - Ages 45-64	▼ LOWER than average		90 %	(89% - 90%)	4,145
Adult Access to Preventive/Ambulatory Care - Ages 65+	▼ LOWER than average		83 %	(79% - 87%)	354
Child and Adolescent Access to Primary Care - Ages 12-24 Months	▼ LOWER than average		77 %	(73% - 81%)	399
Child and Adolescent Access to Primary Care - Ages 2-6 Years	▼ LOWER than average		57 %	(55% - 60%)	1,872
Child and Adolescent Access to Primary Care - Ages 7-11 Years	▼ LOWER than average		55 %	(52% - 57%)	1,638
Child and Adolescent Access to Primary Care - Ages 12-19 Years	▼ LOWER than average		55 %	(53% - 57%)	2,549
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	■ AVERAGE		93 %	(90% - 95%)	434
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	■ AVERAGE		86 %	(82% - 89%)	354
<b>Asthma</b>					
Use of Appropriate Medication	■ AVERAGE		84 %	(78% - 88%)	228
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	▼ LOWER than average		81 %	(78% - 84%)	661
Eye Exam	▼ LOWER than average		48 %	(44% - 51%)	661
Cholesterol Test (LDL-C or Bad Cholesterol)	▼ LOWER than average		71 %	(67% - 74%)	661
Kidney Disease Screening	▼ LOWER than average		73 %	(69% - 76%)	661

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.

## Clark County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	▼ LOWER than average		75 %	(74% - 76%)	6,842
Adult Access to Preventive/Ambulatory Care - Ages 45-64	▼ LOWER than average		84 %	(83% - 85%)	5,914
Adult Access to Preventive/Ambulatory Care - Ages 65+	▼ LOWER than average		72 %	(68% - 75%)	638
Child and Adolescent Access to Primary Care - Ages 12-24 Months	▼ LOWER than average		61 %	(56% - 66%)	367
Child and Adolescent Access to Primary Care - Ages 2-6 Years	▼ LOWER than average		41 %	(39% - 43%)	2,382
Child and Adolescent Access to Primary Care - Ages 7-11 Years	▼ LOWER than average		43 %	(41% - 45%)	2,264
Child and Adolescent Access to Primary Care - Ages 12-19 Years	▼ LOWER than average		46 %	(44% - 47%)	3,787
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	■ AVERAGE		92 %	(89% - 94%)	557
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	■ AVERAGE		88 %	(85% - 91%)	602
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	■ AVERAGE		20 %	(15% - 26%)	212
<b>Asthma</b>					
Use of Appropriate Medication	● BETTER than average		93 %	(90% - 95%)	388
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	■ AVERAGE		84 %	(82% - 86%)	1,278
Eye Exam	▼ LOWER than average		47 %	(45% - 50%)	1,278
Cholesterol Test (LDL-C or Bad Cholesterol)	▼ LOWER than average		70 %	(67% - 72%)	1,278

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP






**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.






## Diabetes





## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Columbia County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: gray;">■</span> AVERAGE		88 %	(83% - 92%)	184
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: gray;">■</span> AVERAGE		91 %	(87% - 94%)	278
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: gray;">■</span> AVERAGE		77 %	(71% - 82%)	208
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: gray;">■</span> AVERAGE		76 %	(73% - 79%)	803
Antidepressants	<span style="color: green;">●</span> BETTER than average		95 %	(93% - 96%)	1,354
Antacid Medication (Proton Pump Inhibitors)	<span style="color: green;">●</span> BETTER than average		95 %	(93% - 96%)	768
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	<span style="color: red;">▼</span> LOWER than average		85 %	(83% - 88%)	901
Medication for attention deficit hyperactivity disorder	<span style="color: red;">▼</span> LOWER than average		54 %	(50% - 59%)	445
<b>Health Screenings</b>					
Screening for Cervical Cancer	<span style="color: gray;">■</span> AVERAGE		65 %	(59% - 71%)	219
Adolescent Well-Care Visits	<span style="color: gray;">■</span> AVERAGE		29 %	(24% - 34%)	277

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Cowlitz County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	▼ LOWER than average		76 %	(75% - 78%)	3,109
Adult Access to Preventive/Ambulatory Care - Ages 45-64	▼ LOWER than average		89 %	(87% - 90%)	2,805
Adult Access to Preventive/Ambulatory Care - Ages 65+	▼ LOWER than average		79 %	(73% - 85%)	178
Child and Adolescent Access to Primary Care - Ages 12-24 Months	■ AVERAGE		86 %	(82% - 89%)	330
Child and Adolescent Access to Primary Care - Ages 2-6 Years	▼ LOWER than average		67 %	(65% - 69%)	1,740
Child and Adolescent Access to Primary Care - Ages 7-11 Years	▼ LOWER than average		62 %	(59% - 64%)	1,662
Child and Adolescent Access to Primary Care - Ages 12-19 Years	▼ LOWER than average		61 %	(59% - 63%)	2,463
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	● BETTER than average		96 %	(93% - 97%)	424
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	■ AVERAGE		87 %	(82% - 90%)	334
<b>Asthma</b>					
Use of Appropriate Medication	■ AVERAGE		88 %	(84% - 92%)	254
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	▼ LOWER than average		82 %	(79% - 85%)	698
Eye Exam	▼ LOWER than average		49 %	(45% - 52%)	698
Cholesterol Test (LDL-C or Bad Cholesterol)	■ AVERAGE		75 %	(71% - 78%)	698
Kidney Disease Screening	■ AVERAGE		78 %	(75% - 81%)	698

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.






# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Douglas County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		84 %	(82% - 85%)	1,747
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: gray;">■</span> AVERAGE		90 %	(88% - 92%)	1,344
Child and Adolescent Access to Primary Care - Ages 12–24 Months	<span style="color: green;">●</span> BETTER than average		95 %	(92% - 97%)	303
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: gray;">■</span> AVERAGE		77 %	(75% - 79%)	1,507
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: green;">●</span> BETTER than average		80 %	(78% - 82%)	1,305
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: gray;">■</span> AVERAGE		77 %	(75% - 79%)	1,877
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> AVERAGE		94 %	(91% - 95%)	471
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> AVERAGE		86 %	(81% - 90%)	197
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		89 %	(84% - 92%)	239
Eye Exam	<span style="color: gray;">■</span> AVERAGE		58 %	(52% - 64%)	239
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> AVERAGE		74 %	(68% - 79%)	239
Kidney Disease Screening	<span style="color: gray;">■</span> AVERAGE		79 %	(73% - 84%)	239
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: red;">▼</span> LOWER than average		74 %	(73% - 76%)	4,058
Antidepressants	<span style="color: gray;">■</span> AVERAGE		90 %	(89% - 90%)	7,273

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.

## Ferry County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	▼ LOWER than average		71 %	(65% - 77%)	225
Adult Access to Preventive/Ambulatory Care - Ages 45–64	▼ LOWER than average		79 %	(74% - 83%)	274
Child and Adolescent Access to Primary Care - Ages 2–6 Years	▼ LOWER than average		43 %	(36% - 50%)	176
Child and Adolescent Access to Primary Care - Ages 12–19 Years	▼ LOWER than average		65 %	(58% - 71%)	199
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	▼ LOWER than average		69 %	(65% - 72%)	710
Antidepressants	■ AVERAGE		88 %	(87% - 90%)	1,462
Antacid Medication (Proton Pump Inhibitors)	■ AVERAGE		86 %	(84% - 89%)	709
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	▼ LOWER than average		82 %	(79% - 84%)	1,179
Medication for attention deficit hyperactivity disorder	■ AVERAGE		64 %	(59% - 69%)	343
<b>Health Screenings</b>					
Screening for Cervical Cancer	▼ LOWER than average		55 %	(49% - 61%)	247
Adolescent Well-Care Visits	▼ LOWER than average		20 %	(15% - 25%)	268

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Franklin County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> <b>BETTER than average</b>		89 %	(88% - 90%)	4,266
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: green;">●</span> <b>BETTER than average</b>		95 %	(94% - 95%)	2,873
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> <b>AVERAGE</b>		85 %	(81% - 89%)	259
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: gray;">■</span> <b>AVERAGE</b>		92 %	(90% - 94%)	785
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		79 %	(77% - 80%)	4,069
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		79 %	(78% - 81%)	3,425
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		75 %	(74% - 76%)	4,340
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: red;">▼</span> <b>LOWER than average</b>		85 %	(83% - 87%)	1,207
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> <b>AVERAGE</b>		86 %	(82% - 89%)	411
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(83% - 91%)	256
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: red;">▼</span> <b>LOWER than average</b>		83 %	(80% - 85%)	828
Eye Exam	<span style="color: gray;">■</span> <b>AVERAGE</b>		55 %	(51% - 58%)	828
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> <b>AVERAGE</b>		74 %	(71% - 77%)	828
Kidney Disease Screening	<span style="color: red;">▼</span> <b>LOWER than average</b>		74 %	(71% - 77%)	828

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Garfield County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Scripts*
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: gray;">■</span> AVERAGE		76 %	(73% - 79%)	672
Antidepressants	<span style="color: red;">▼</span> LOWER than average		83 %	(80% - 86%)	616
Antacid Medication (Proton Pump Inhibitors)	<span style="color: gray;">■</span> AVERAGE		92 %	(89% - 94%)	395
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	<span style="color: green;">●</span> BETTER than average		94 %	(92% - 95%)	747

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Grant County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		85 %	(84% - 86%)	3,891
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: gray;">■</span> AVERAGE		91 %	(90% - 92%)	2,866
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> AVERAGE		86 %	(81% - 90%)	234
Child and Adolescent Access to Primary Care - Ages 12–24 Months	<span style="color: gray;">■</span> AVERAGE		87 %	(85% - 90%)	567
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: gray;">■</span> AVERAGE		74 %	(73% - 76%)	2,868
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: gray;">■</span> AVERAGE		75 %	(73% - 77%)	2,542
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: red;">▼</span> LOWER than average		71 %	(70% - 73%)	3,803
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> AVERAGE		91 %	(90% - 93%)	1,244
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> AVERAGE		88 %	(84% - 90%)	471
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: gray;">■</span> AVERAGE		29 %	(23% - 37%)	160
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> AVERAGE		88 %	(83% - 91%)	272
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		85 %	(82% - 88%)	706
Eye Exam	<span style="color: red;">▼</span> LOWER than average		46 %	(43% - 50%)	706
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> AVERAGE		73 %	(69% - 76%)	706

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.


Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.








**LEGEND** ● **BETTER** than average Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER** than average Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.






## Diabetes



## Generic Prescription Drugs



## Health Screenings





# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Grays Harbor County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	▼ LOWER than average		83 %	(81% - 84%)	4,181
Adult Access to Preventive/Ambulatory Care - Ages 45-64	■ AVERAGE		91 %	(90% - 92%)	4,498
Adult Access to Preventive/Ambulatory Care - Ages 65+	■ AVERAGE		88 %	(85% - 91%)	404
Child and Adolescent Access to Primary Care - Ages 12-24 Months	▼ LOWER than average		84 %	(80% - 87%)	404
Child and Adolescent Access to Primary Care - Ages 2-6 Years	▼ LOWER than average		68 %	(66% - 70%)	1,906
Child and Adolescent Access to Primary Care - Ages 7-11 Years	▼ LOWER than average		67 %	(65% - 69%)	1,733
Child and Adolescent Access to Primary Care - Ages 12-19 Years	▼ LOWER than average		68 %	(66% - 70%)	2,711
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	▼ LOWER than average		86 %	(82% - 89%)	385
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	■ AVERAGE		86 %	(82% - 89%)	385
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	■ AVERAGE		24 %	(19% - 30%)	227
<b>Asthma</b>					
Use of Appropriate Medication	▼ LOWER than average		79 %	(74% - 84%)	257
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	▼ LOWER than average		81 %	(78% - 83%)	902
Eye Exam	▼ LOWER than average		54 %	(50% - 57%)	902
Cholesterol Test (LDL-C or Bad Cholesterol)	▼ LOWER than average		67 %	(64% - 70%)	902

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP





**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.






## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

## LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average

Must have 100+ patients per measure to be publicly reported.

## Island County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	AVERAGE		86 %	(85% - 88%)	2,319
Adult Access to Preventive/Ambulatory Care - Ages 45-64	AVERAGE		93 %	(92% - 94%)	4,213
Adult Access to Preventive/Ambulatory Care - Ages 65+	AVERAGE		87 %	(83% - 90%)	408
Child and Adolescent Access to Primary Care - Ages 2-6 Years	LOWER than average		67 %	(63% - 71%)	574
Child and Adolescent Access to Primary Care - Ages 7-11 Years	LOWER than average		70 %	(67% - 74%)	603
Child and Adolescent Access to Primary Care - Ages 12-19 Years	AVERAGE		74 %	(71% - 76%)	1,317
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	AVERAGE		90 %	(85% - 93%)	194
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	AVERAGE		91 %	(86% - 94%)	253
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	BETTER than average		48 %	(41% - 55%)	193
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	AVERAGE		85 %	(82% - 88%)	643
Eye Exam	AVERAGE		59 %	(55% - 63%)	643
Cholesterol Test (LDL-C or Bad Cholesterol)	AVERAGE		77 %	(73% - 80%)	643
Kidney Disease Screening	LOWER than average		74 %	(71% - 77%)	643
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	AVERAGE		77 %	(76% - 78%)	10,676

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.






# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.


## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Jefferson County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	▼ LOWER than average		81 %	(79% - 83%)	1,282
Adult Access to Preventive/Ambulatory Care - Ages 45–64	■ AVERAGE		92 %	(91% - 93%)	1,942
Adult Access to Preventive/Ambulatory Care - Ages 65+	■ AVERAGE		84 %	(79% - 88%)	237
Child and Adolescent Access to Primary Care - Ages 2–6 Years	▼ LOWER than average		26 %	(22% - 29%)	602
Child and Adolescent Access to Primary Care - Ages 7–11 Years	▼ LOWER than average		43 %	(38% - 48%)	333
Child and Adolescent Access to Primary Care - Ages 12–19 Years	▼ LOWER than average		49 %	(45% - 53%)	626
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	■ AVERAGE		88 %	(83% - 91%)	249
Eye Exam	▼ LOWER than average		49 %	(43% - 55%)	249
Cholesterol Test (LDL-C or Bad Cholesterol)	■ AVERAGE		76 %	(70% - 81%)	249
Kidney Disease Screening	■ AVERAGE		80 %	(74% - 84%)	249
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	■ AVERAGE		77 %	(75% - 78%)	4,264
Antidepressants	▼ LOWER than average		85 %	(84% - 86%)	7,234
Antacid Medication (Proton Pump Inhibitors)	▼ LOWER than average		86 %	(84% - 87%)	2,276
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	■ AVERAGE		88 %	(87% - 89%)	5,071

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP






**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## King County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> <b>BETTER than average</b>		86 %	(86% - 86%)	119,314
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: gray;">■</span> <b>AVERAGE</b>		92 %	(92% - 92%)	123,093
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> <b>AVERAGE</b>		90 %	(89% - 91%)	11,861
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: green;">●</span> <b>BETTER than average</b>		91 %	(91% - 92%)	9,100
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		79 %	(78% - 79%)	43,835
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		80 %	(79% - 80%)	35,371
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		78 %	(77% - 78%)	55,541
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: green;">●</span> <b>BETTER than average</b>		95 %	(94% - 95%)	12,067
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> <b>AVERAGE</b>		87 %	(86% - 88%)	10,566
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: green;">●</span> <b>BETTER than average</b>		28 %	(26% - 29%)	4,651
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: green;">●</span> <b>BETTER than average</b>		89 %	(88% - 90%)	4,285
<b>COPD</b>					
Use of Spirometry Testing in the Assessment and Diagnosis of COPD	<span style="color: green;">●</span> <b>BETTER than average</b>		48 %	(44% - 52%)	651
<b>Depression</b>					
Antidepressant Medication (12 Weeks)	<span style="color: gray;">■</span> <b>AVERAGE</b>		68 %	(66% - 69%)	3,689

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Depression




## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



## LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average


Must have 100+ patients per measure to be publicly reported.

## Heart Disease

Cholesterol-Lowering Medication



**AVERAGE**




71 % (70% - 73%)

4,652

Beta Blockers



**AVERAGE**



80 % (74% - 85%)

187

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Kitsap County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> <b>BETTER than average</b>		87 %	(86% - 87%)	13,156
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: green;">●</span> <b>BETTER than average</b>		94 %	(94% - 94%)	17,493
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: green;">●</span> <b>BETTER than average</b>		92 %	(90% - 94%)	1,146
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(86% - 90%)	912
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		74 %	(73% - 76%)	4,796
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		75 %	(74% - 77%)	4,515
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		77 %	(76% - 78%)	7,711
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> <b>AVERAGE</b>		93 %	(91% - 94%)	1,269
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: red;">▼</span> <b>LOWER than average</b>		83 %	(81% - 85%)	1,216
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: gray;">■</span> <b>AVERAGE</b>		27 %	(24% - 30%)	865
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> <b>AVERAGE</b>		85 %	(82% - 88%)	628
<b>Depression</b>					
Antidepressant Medication (12 Weeks)	<span style="color: gray;">■</span> <b>AVERAGE</b>		67 %	(63% - 70%)	573
Antidepressant Medication (6 Months)	<span style="color: gray;">■</span> <b>AVERAGE</b>		52 %	(48% - 56%)	573

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Kittitas County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		86 %	(85% - 88%)	2,154
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: green;">●</span> BETTER than average		95 %	(94% - 96%)	2,336
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> AVERAGE		92 %	(87% - 95%)	180
Child and Adolescent Access to Primary Care - Ages 12–24 Months	<span style="color: gray;">■</span> AVERAGE		92 %	(88% - 95%)	230
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: gray;">■</span> AVERAGE		77 %	(74% - 79%)	1,013
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: gray;">■</span> AVERAGE		78 %	(75% - 80%)	891
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: gray;">■</span> AVERAGE		74 %	(72% - 76%)	1,381
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> AVERAGE		93 %	(89% - 96%)	215
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> AVERAGE		90 %	(85% - 93%)	211
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		84 %	(80% - 88%)	322
Eye Exam	<span style="color: gray;">■</span> AVERAGE		61 %	(56% - 67%)	322
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> AVERAGE		74 %	(69% - 78%)	322
Kidney Disease Screening	<span style="color: gray;">■</span> AVERAGE		76 %	(71% - 80%)	322
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: green;">●</span> BETTER than average		79 %	(78% - 80%)	6,615

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.

## Klickitat County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	▼ LOWER than average		69 %	(65% - 72%)	610
Adult Access to Preventive/Ambulatory Care - Ages 45–64	▼ LOWER than average		78 %	(75% - 81%)	600
Child and Adolescent Access to Primary Care - Ages 2–6 Years	▼ LOWER than average		31 %	(25% - 36%)	272
Child and Adolescent Access to Primary Care - Ages 7–11 Years	▼ LOWER than average		30 %	(25% - 36%)	257
Child and Adolescent Access to Primary Care - Ages 12–19 Years	▼ LOWER than average		38 %	(33% - 43%)	396
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	● BETTER than average		81 %	(79% - 83%)	1,825
Antidepressants	■ AVERAGE		88 %	(87% - 89%)	3,837
Antacid Medication (Proton Pump Inhibitors)	■ AVERAGE		88 %	(86% - 89%)	1,565
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	● BETTER than average		92 %	(91% - 93%)	2,046
Medication for attention deficit hyperactivity disorder	■ AVERAGE		65 %	(62% - 68%)	1,242
<b>Health Screenings</b>					
Screening for Breast Cancer - Ages 52–69	▼ LOWER than average		52 %	(46% - 57%)	277
Screening for Colon Cancer	▼ LOWER than average		38 %	(32% - 44%)	232
Screening for Cervical Cancer	▼ LOWER than average		53 %	(49% - 56%)	670
Adolescent Well-Care Visits	▼ LOWER than average		21 %	(18% - 24%)	561

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





## LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average


Significantly lower  
than regional average

Must have 160+ patients per measure to be publicly reported.

## Lewis County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.






# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Lincoln County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		84 %	(81% - 87%)	520
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: gray;">■</span> AVERAGE		93 %	(91% - 95%)	685
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: gray;">■</span> AVERAGE		77 %	(72% - 82%)	304
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: green;">●</span> BETTER than average		82 %	(77% - 86%)	300
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: green;">●</span> BETTER than average		80 %	(76% - 83%)	478
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: red;">▼</span> LOWER than average		74 %	(72% - 76%)	2,078
Antidepressants	<span style="color: red;">▼</span> LOWER than average		84 %	(83% - 86%)	2,714
Antacid Medication (Proton Pump Inhibitors)	<span style="color: red;">▼</span> LOWER than average		83 %	(80% - 85%)	1,153
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	<span style="color: red;">▼</span> LOWER than average		86 %	(85% - 88%)	2,756
Medication for attention deficit hyperactivity disorder	<span style="color: red;">▼</span> LOWER than average		54 %	(50% - 58%)	690
<b>Health Screenings</b>					
Screening for Breast Cancer - Ages 52–69	<span style="color: gray;">■</span> AVERAGE		69 %	(64% - 74%)	303
Screening for Colon Cancer	<span style="color: gray;">■</span> AVERAGE		48 %	(43% - 54%)	317
Screening for Cervical Cancer	<span style="color: gray;">■</span> AVERAGE		70 %	(67% - 74%)	603
Adolescent Well-Care Visits	<span style="color: gray;">■</span> AVERAGE		33 %	(29% - 37%)	625

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.

















**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.

## Mason County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: gray;">■</span> AVERAGE		85 %	(84% - 86%)	2,985
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: gray;">■</span> AVERAGE		92 %	(91% - 93%)	4,054
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: green;">●</span> BETTER than average		94 %	(90% - 96%)	236
Child and Adolescent Access to Primary Care - Ages 12–24 Months	<span style="color: red;">▼</span> LOWER than average		83 %	(78% - 87%)	247
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: red;">▼</span> LOWER than average		61 %	(58% - 64%)	1,144
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: red;">▼</span> LOWER than average		61 %	(58% - 64%)	1,043
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: red;">▼</span> LOWER than average		61 %	(59% - 63%)	1,667
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: green;">●</span> BETTER than average		96 %	(93% - 98%)	272
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> AVERAGE		85 %	(80% - 89%)	293
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: green;">●</span> BETTER than average		33 %	(27% - 40%)	189
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		86 %	(83% - 88%)	726
Eye Exam	<span style="color: gray;">■</span> AVERAGE		59 %	(56% - 63%)	726
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> AVERAGE		75 %	(71% - 78%)	726
Kidney Disease Screening	<span style="color: gray;">■</span> AVERAGE		81 %	(78% - 83%)	726

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.


## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Okanogan County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	▼ LOWER than average		83 %	(81% - 84%)	1,952
Adult Access to Preventive/Ambulatory Care - Ages 45-64	▼ LOWER than average		90 %	(88% - 91%)	1,825
Child and Adolescent Access to Primary Care - Ages 12-24 Months	▼ LOWER than average		83 %	(79% - 87%)	378
Child and Adolescent Access to Primary Care - Ages 2-6 Years	▼ LOWER than average		67 %	(65% - 69%)	1,740
Child and Adolescent Access to Primary Care - Ages 7-11 Years	▼ LOWER than average		67 %	(64% - 69%)	1,395
Child and Adolescent Access to Primary Care - Ages 12-19 Years	▼ LOWER than average		69 %	(67% - 71%)	1,823
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	■ AVERAGE		91 %	(87% - 93%)	405
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	■ AVERAGE		87 %	(81% - 91%)	162
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	■ AVERAGE		84 %	(79% - 87%)	339
Eye Exam	● BETTER than average		67 %	(62% - 72%)	339
Cholesterol Test (LDL-C or Bad Cholesterol)	■ AVERAGE		75 %	(70% - 79%)	339
Kidney Disease Screening	■ AVERAGE		76 %	(71% - 80%)	339
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	▼ LOWER than average		60 %	(59% - 62%)	6,167
Antidepressants	● BETTER than average		90 %	(90% - 91%)	9,041

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.


Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.

## Pacific County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	▼ LOWER than average		82 %	(79% - 84%)	961
Adult Access to Preventive/Ambulatory Care - Ages 45–64	■ AVERAGE		91 %	(89% - 93%)	1,209
Child and Adolescent Access to Primary Care - Ages 2–6 Years	▼ LOWER than average		64 %	(59% - 68%)	462
Child and Adolescent Access to Primary Care - Ages 7–11 Years	▼ LOWER than average		62 %	(57% - 66%)	442
Child and Adolescent Access to Primary Care - Ages 12–19 Years	▼ LOWER than average		63 %	(59% - 67%)	632
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	▼ LOWER than average		81 %	(76% - 85%)	271
Eye Exam	■ AVERAGE		57 %	(51% - 63%)	271
Cholesterol Test (LDL-C or Bad Cholesterol)	■ AVERAGE		72 %	(66% - 77%)	271
Kidney Disease Screening	■ AVERAGE		75 %	(69% - 80%)	271
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	■ AVERAGE		78 %	(77% - 79%)	4,523
Antidepressants	■ AVERAGE		88 %	(88% - 89%)	5,481
Antacid Medication (Proton Pump Inhibitors)	▼ LOWER than average		87 %	(85% - 88%)	2,479
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	■ AVERAGE		88 %	(88% - 89%)	5,590
Medication for attention deficit hyperactivity disorder	▼ LOWER than average		62 %	(60% - 65%)	1,214

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

## LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average




**LOWER**  
than average

Significantly lower  
than regional average

Must have 160+ patients per measure to be publicly reported.

## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.

















**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Pend Oreille County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	▼ LOWER than average		78 %	(74% - 81%)	423
Adult Access to Preventive/Ambulatory Care - Ages 45–64	▼ LOWER than average		81 %	(77% - 84%)	566
Child and Adolescent Access to Primary Care - Ages 2–6 Years	▼ LOWER than average		65 %	(59% - 70%)	275
Child and Adolescent Access to Primary Care - Ages 7–11 Years	▼ LOWER than average		64 %	(57% - 70%)	217
Child and Adolescent Access to Primary Care - Ages 12–19 Years	■ AVERAGE		71 %	(66% - 75%)	381
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	▼ LOWER than average		71 %	(69% - 74%)	1,523
Antidepressants	▼ LOWER than average		84 %	(82% - 85%)	3,029
Antacid Medication (Proton Pump Inhibitors)	■ AVERAGE		87 %	(85% - 89%)	1,414
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	■ AVERAGE		90 %	(89% - 91%)	1,803
Medication for attention deficit hyperactivity disorder	■ AVERAGE		68 %	(65% - 72%)	716
<b>Health Screenings</b>					
Screening for Breast Cancer - Ages 52–69	▼ LOWER than average		54 %	(47% - 60%)	252
Screening for Colon Cancer	■ AVERAGE		44 %	(38% - 50%)	237
Screening for Cervical Cancer	▼ LOWER than average		58 %	(53% - 62%)	507
Adolescent Well-Care Visits	▼ LOWER than average		22 %	(19% - 26%)	479

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**


- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Pierce County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.


## Depression




## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average


Must have 100+ patients per measure to be publicly reported.

## Heart Disease

Cholesterol-Lowering Medication



**AVERAGE**



73 % (71% - 75%)

2,182





**LEGEND**













- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.

## San Juan County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> <b>BETTER than average</b>		92 %	(89% - 94%)	375
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: gray;">■</span> <b>AVERAGE</b>		94 %	(92% - 95%)	898
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		83 %	(78% - 86%)	280
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: green;">●</span> <b>BETTER than average</b>		80 %	(78% - 81%)	2,326
Antidepressants	<span style="color: gray;">■</span> <b>AVERAGE</b>		89 %	(88% - 90%)	3,218
Antacid Medication (Proton Pump Inhibitors)	<span style="color: red;">▼</span> <b>LOWER than average</b>		86 %	(84% - 88%)	1,136
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	<span style="color: gray;">■</span> <b>AVERAGE</b>		90 %	(89% - 91%)	2,407
Medication for attention deficit hyperactivity disorder	<span style="color: red;">▼</span> <b>LOWER than average</b>		60 %	(56% - 64%)	606
<b>Health Screenings</b>					
Screening for Breast Cancer - Ages 52-69	<span style="color: gray;">■</span> <b>AVERAGE</b>		63 %	(59% - 67%)	591
Screening for Colon Cancer	<span style="color: red;">▼</span> <b>LOWER than average</b>		43 %	(38% - 47%)	482
Screening for Cervical Cancer	<span style="color: gray;">■</span> <b>AVERAGE</b>		68 %	(64% - 71%)	655
Adolescent Well-Care Visits	<span style="color: gray;">■</span> <b>AVERAGE</b>		28 %	(24% - 32%)	454

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Skagit County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER** than average Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER** than average Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.














**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Skamania County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	▼ LOWER than average		68 %	(63% - 74%)	278
Adult Access to Preventive/Ambulatory Care - Ages 45–64	▼ LOWER than average		76 %	(70% - 82%)	193
Child and Adolescent Access to Primary Care - Ages 2–6 Years	▼ LOWER than average		50 %	(43% - 56%)	242
Child and Adolescent Access to Primary Care - Ages 12–19 Years	▼ LOWER than average		38 %	(31% - 46%)	160
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	▼ LOWER than average		69 %	(65% - 73%)	644
Antidepressants	▼ LOWER than average		80 %	(78% - 82%)	1,657
Antacid Medication (Proton Pump Inhibitors)	▼ LOWER than average		76 %	(73% - 80%)	580
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	● BETTER than average		95 %	(94% - 97%)	814
Medication for attention deficit hyperactivity disorder	■ AVERAGE		63 %	(59% - 66%)	739
<b>Health Screenings</b>					
Screening for Cervical Cancer	▼ LOWER than average		56 %	(50% - 62%)	264
Adolescent Well-Care Visits	▼ LOWER than average		17 %	(14% - 21%)	370

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Snohomish County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> <b>BETTER than average</b>		87 %	(86% - 87%)	41,417
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: gray;">■</span> <b>AVERAGE</b>		92 %	(92% - 93%)	42,060
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> <b>AVERAGE</b>		90 %	(89% - 91%)	3,414
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: gray;">■</span> <b>AVERAGE</b>		91 %	(90% - 92%)	3,431
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		79 %	(78% - 79%)	17,568
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		80 %	(79% - 81%)	14,500
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		78 %	(77% - 78%)	22,447
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> <b>AVERAGE</b>		92 %	(91% - 93%)	4,589
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(86% - 89%)	3,991
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: gray;">■</span> <b>AVERAGE</b>		23 %	(21% - 25%)	1,924
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> <b>AVERAGE</b>		85 %	(83% - 87%)	1,667
<b>COPD</b>					
Use of Spirometry Testing in the Assessment and Diagnosis of COPD	<span style="color: gray;">■</span> <b>AVERAGE</b>		37 %	(31% - 43%)	226
<b>Depression</b>					
Antidepressant Medication (12 Weeks)	<span style="color: gray;">■</span> <b>AVERAGE</b>		64 %	(61% - 66%)	1,433

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.







**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Depression




## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average


Must have 100+ patients per measure to be publicly reported.

## Heart Disease

Cholesterol-Lowering Medication



**AVERAGE**



70 % (68% - 72%)

1,725



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.

## Spokane County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: gray;">■</span> AVERAGE		84 %	(84% - 85%)	26,589
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: red;">▼</span> LOWER than average		90 %	(90% - 91%)	25,733
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: red;">▼</span> LOWER than average		82 %	(80% - 83%)	2,057
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: gray;">■</span> AVERAGE		91 %	(90% - 92%)	3,063
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: gray;">■</span> AVERAGE		76 %	(75% - 76%)	15,281
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: gray;">■</span> AVERAGE		77 %	(76% - 78%)	12,516
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: green;">●</span> BETTER than average		77 %	(76% - 78%)	18,156
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: red;">▼</span> LOWER than average		88 %	(87% - 89%)	3,784
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: red;">▼</span> LOWER than average		84 %	(83% - 85%)	2,543
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: red;">▼</span> LOWER than average		16 %	(14% - 18%)	1,410
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> AVERAGE		84 %	(82% - 86%)	1,660
<b>COPD</b>					
Use of Spirometry Testing in the Assessment and Diagnosis of COPD	<span style="color: gray;">■</span> AVERAGE		36 %	(30% - 42%)	229
<b>Depression</b>					
Antidepressant Medication (12 Weeks)	<span style="color: gray;">■</span> AVERAGE		68 %	(65% - 71%)	906

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP





**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 180+ patients per measure to be publicly reported.





## Depression








## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average


Must have 160+ patients per measure to be publicly reported.

## Heart Disease

Cholesterol-Lowering Medication



**LOWER**  
than average



**66 % (63 % - 69 %)**

**1,098**



## LEGEND



**BETTER**  
than average

Significantly better  
than regional average



**AVERAGE**

No significant difference  
from regional average



**LOWER**  
than average

Significantly lower  
than regional average

Must have 180+ patients per measure to be publicly reported.

## Stevens County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	LOWER than average		82 %	(80% - 84%)	1,699
Adult Access to Preventive/Ambulatory Care - Ages 45-64	LOWER than average		87 %	(86% - 89%)	2,145
Child and Adolescent Access to Primary Care - Ages 12-24 Months	LOWER than average		79 %	(72% - 84%)	163
Child and Adolescent Access to Primary Care - Ages 2-6 Years	LOWER than average		63 %	(60% - 66%)	891
Child and Adolescent Access to Primary Care - Ages 7-11 Years	LOWER than average		65 %	(61% - 68%)	866
Child and Adolescent Access to Primary Care - Ages 12-19 Years	AVERAGE		72 %	(69% - 74%)	1,352
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	LOWER than average		81 %	(76% - 85%)	243
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	AVERAGE		85 %	(79% - 90%)	166
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	AVERAGE		83 %	(78% - 86%)	374
Eye Exam	LOWER than average		50 %	(45% - 55%)	374
Cholesterol Test (LDL-C or Bad Cholesterol)	LOWER than average		69 %	(64% - 73%)	374
Kidney Disease Screening	AVERAGE		75 %	(70% - 79%)	374
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	BETTER than average		79 %	(78% - 80%)	4,831
Antidepressants	AVERAGE		89 %	(88% - 90%)	9,604

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.





**LEGEND**














- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Thurston County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: green;">●</span> <b>BETTER than average</b>		88 %	(87% - 88%)	16,072
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: green;">●</span> <b>BETTER than average</b>		94 %	(94% - 94%)	19,081
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: green;">●</span> <b>BETTER than average</b>		92 %	(91% - 93%)	1,553
Child and Adolescent Access to Primary Care - Ages 12–24 Months	<span style="color: gray;">■</span> <b>AVERAGE</b>		91 %	(89% - 92%)	1,188
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		76 %	(75% - 77%)	5,928
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		77 %	(76% - 79%)	5,375
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: green;">●</span> <b>BETTER than average</b>		78 %	(77% - 79%)	8,894
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> <b>AVERAGE</b>		92 %	(91% - 94%)	1,615
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> <b>AVERAGE</b>		86 %	(85% - 88%)	1,668
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: green;">●</span> <b>BETTER than average</b>		33 %	(30% - 37%)	830
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> <b>AVERAGE</b>		86 %	(83% - 88%)	721
<b>Depression</b>					
Antidepressant Medication (12 Weeks)	<span style="color: green;">●</span> <b>BETTER than average</b>		71 %	(68% - 74%)	723
Antidepressant Medication (6 Months)	<span style="color: green;">●</span> <b>BETTER than average</b>		56 %	(53% - 60%)	723

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 100+ patients per measure to be publicly reported.

## Wahkiakum County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Scripts*
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: green;">●</span> <b>BETTER than average</b>		82 %	(78% - 84%)	612
Antidepressants	<span style="color: gray;">■</span> <b>AVERAGE</b>		91 %	(89% - 93%)	1,000
Antacid Medication (Proton Pump Inhibitors)	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(84% - 91%)	384
Medication for ACE inhibitors or angiotensin II receptor blockers (ARBs) used to treat High Blood Pressure	<span style="color: green;">●</span> <b>BETTER than average</b>		92 %	(90% - 94%)	739
Medication for attention deficit hyperactivity disorder	<span style="color: gray;">■</span> <b>AVERAGE</b>		66 %	(61% - 71%)	342

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**















- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Walla Walla County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> BETTER than average		88 %	(87% - 89%)	3,154
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: gray;">■</span> AVERAGE		93 %	(92% - 94%)	3,182
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> AVERAGE		92 %	(88% - 95%)	205
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: gray;">■</span> AVERAGE		91 %	(88% - 94%)	354
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: green;">●</span> BETTER than average		79 %	(77% - 80%)	1,871
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: green;">●</span> BETTER than average		81 %	(79% - 83%)	1,795
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: green;">●</span> BETTER than average		79 %	(77% - 80%)	2,570
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> AVERAGE		90 %	(88% - 92%)	610
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> AVERAGE		86 %	(82% - 89%)	405
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: gray;">■</span> AVERAGE		24 %	(18% - 31%)	175
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> AVERAGE		85 %	(80% - 89%)	223
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		85 %	(83% - 88%)	641
Eye Exam	<span style="color: gray;">■</span> AVERAGE		60 %	(56% - 63%)	641
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: gray;">■</span> AVERAGE		77 %	(74% - 80%)	641

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP






**LEGEND** ● **BETTER** than average Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER** than average Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.






## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



# COMMUNITY CHECKUP

**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Whatcom County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20-44	<span style="color: green;">●</span> <b>BETTER than average</b>		87 %	(86% - 87%)	12,227
Adult Access to Preventive/Ambulatory Care - Ages 45-64	<span style="color: green;">●</span> <b>BETTER than average</b>		94 %	(93% - 94%)	12,326
Adult Access to Preventive/Ambulatory Care - Ages 65+	<span style="color: gray;">■</span> <b>AVERAGE</b>		90 %	(88% - 92%)	839
Child and Adolescent Access to Primary Care - Ages 12-24 Months	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(86% - 90%)	1,220
Child and Adolescent Access to Primary Care - Ages 2-6 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		75 %	(74% - 77%)	5,788
Child and Adolescent Access to Primary Care - Ages 7-11 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		74 %	(73% - 75%)	4,780
Child and Adolescent Access to Primary Care - Ages 12-19 Years	<span style="color: gray;">■</span> <b>AVERAGE</b>		75 %	(74% - 76%)	7,434
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> <b>AVERAGE</b>		93 %	(91% - 94%)	1,459
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> <b>AVERAGE</b>		88 %	(86% - 90%)	1,100
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	<span style="color: red;">▼</span> <b>LOWER than average</b>		17 %	(14% - 20%)	633
<b>Asthma</b>					
Use of Appropriate Medication	<span style="color: gray;">■</span> <b>AVERAGE</b>		89 %	(87% - 92%)	642
<b>Depression</b>					
Antidepressant Medication (12 Weeks)	<span style="color: gray;">■</span> <b>AVERAGE</b>		67 %	(63% - 72%)	401
Antidepressant Medication (6 Months)	<span style="color: gray;">■</span> <b>AVERAGE</b>		52 %	(47% - 57%)	401

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.






# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**

- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Whitman County

Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code

		Confidence Interval Range & RATE	RATE	Confidence Interval	Patients*
<b>Access to Care</b>					
Adult Access to Preventive/Ambulatory Care - Ages 20–44	<span style="color: green;">●</span> BETTER than average		90 %	(88% - 91%)	2,000
Adult Access to Preventive/Ambulatory Care - Ages 45–64	<span style="color: green;">●</span> BETTER than average		94 %	(93% - 95%)	1,827
Child and Adolescent Access to Primary Care - Ages 12–24 Months	<span style="color: green;">●</span> BETTER than average		94 %	(91% - 97%)	229
Child and Adolescent Access to Primary Care - Ages 2–6 Years	<span style="color: green;">●</span> BETTER than average		84 %	(81% - 86%)	1,003
Child and Adolescent Access to Primary Care - Ages 7–11 Years	<span style="color: green;">●</span> BETTER than average		85 %	(82% - 87%)	763
Child and Adolescent Access to Primary Care - Ages 12–19 Years	<span style="color: green;">●</span> BETTER than average		87 %	(84% - 88%)	1,120
<b>Appropriate Use of Care</b>					
Avoidance of Antibiotics for Common Cold	<span style="color: gray;">■</span> AVERAGE		89 %	(85% - 92%)	387
Avoidance of X-ray, MRI and CT Scan for Low Back Pain	<span style="color: gray;">■</span> AVERAGE		89 %	(84% - 92%)	217
<b>Diabetes</b>					
Blood Sugar (HbA1c) Test	<span style="color: gray;">■</span> AVERAGE		82 %	(76% - 86%)	251
Eye Exam	<span style="color: gray;">■</span> AVERAGE		52 %	(46% - 58%)	251
Cholesterol Test (LDL-C or Bad Cholesterol)	<span style="color: red;">▼</span> LOWER than average		63 %	(56% - 68%)	251
Kidney Disease Screening	<span style="color: gray;">■</span> AVERAGE		76 %	(70% - 81%)	251
<b>Generic Prescription Drugs</b>					
Cholesterol-Lowering Drugs (Statins)	<span style="color: green;">●</span> BETTER than average		80 %	(79% - 81%)	7,164
Antidepressants	<span style="color: gray;">■</span> AVERAGE		88 %	(87% - 89%)	9,788

\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average ■ **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Generic Prescription Drugs



## Health Screenings



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.



**LEGEND**


- **BETTER than average** Significantly better than regional average
- **AVERAGE** No significant difference from regional average
- ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.

## Yakima County

### Performance results among the all payers insured population, organized by county

County measures are attributed using the patient's zip code



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.




# COMMUNITY CHECKUP


**LEGEND** ● **BETTER than average** Significantly better than regional average  **AVERAGE** No significant difference from regional average ▼ **LOWER than average** Significantly lower than regional average

Must have 160+ patients per measure to be publicly reported.


## Diabetes




## Generic Prescription Drugs



## Health Screenings



## Heart Disease



\* The denominator value affects the accuracy of the results. Based on this number, the Confidence Interval denotes the range within which 95% of results are likely to occur.

Based on claims and encounter data with dates of service between 1/1/2004 - 6/30/2012 and the measurement year of 7/1/2011 - 6/30/2012.